


## Adams County

## Adults

## Current Cigarette Smoking (1987-2002)




+Insufficient data (< 50 total observations) to report for Adams County in 1989-1993.

## Current Smokeless Tobacco Use (1999-2001)


	Entire adult population	Males only
Adams County	insufficient data (< 50 total observations)	insufficient data (< 50 total observations)
Washington State	2.8% (95%CI: 2.5,3.2)	5.6% (95%CI: 4.9,6.4)

## Youth



DATA SOURCES: (ADDITIONAL DOCUMENTATION INCLUDED IN APPENDIX A, [WWW.DOH.WA.GOV/TOBACCO/DATA/COUNTY\\_PROFILES/PROFILES\\_2002/APPNDA.PDF](http://WWW.DOH.WA.GOV/TOBACCO/DATA/COUNTY_PROFILES/PROFILES_2002/APPNDA.PDF))  
 ADULT TOBACCO USE – WA. DEPT OF HEALTH, BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM (BRFSS), 2002 PRELIMINARY ESTIMATE BASED ON FIRST SIX MONTHS OF YEAR; YOUTH TOBACCO USE – WA STATE SURVEY OF ADOLESCENT HEALTH BEHAVIORS (WSSAHB); POPULATION ESTIMATES – OFFICE OF FINANCIAL MGMT(OFM), YEAR 2000 CENSUS; MATERNAL SMOKING DURING PREGNANCY – WA BIRTH CERT SYSTEM; LUNG AND BRONCHUS CANCER – WA STATE CANCER REGISTRY; ESTIMATED TOTAL DEATHS – WA DEATH CERT SYSTEM.

## Pregnant Women




### Summary of the problem

- Although adult cigarette smoking has been lower than Washington State as a whole since 1996, it is not significantly different from the state overall. Our most recent survey data showed that about 13.4% of adults in Adams County smoke cigarettes.
- Not enough information was available to describe adult smokeless tobacco use.
- While it appears greater than the state rate for grades 6, 8 and 10, youth smoking in Adams County is not significantly greater than Washington State as a whole. Youth smokeless tobacco use rates are significantly greater for grade 8 than the state as a whole.
- Maternal smoking during pregnancy is consistently and significantly lower than the state as a whole. In recent years, the trend is decreasing less than the state rate. In 2001, Adams County was among the lowest for rate of maternal smoking in the state. About 4.9% of pregnant women in Adams County smoked during pregnancy, compared to 12.6% statewide.

### People Affected in Adams County

- *Total Population: 16,428 - 18 and over: 10,815 (66%); under 18: 5,613 (34%)*
- *Average number of people diagnosed with Lung and Bronchus cancer each year: 3*
- *Infants born annually to mothers who smoke during pregnancy: 10*
- *Estimated total deaths due to tobacco use each year: 20*

Karen Potts, RN  
 Adams County Health District  
 108 West Main  
 Ritzville, WA 99169  
 (509) 659-3320  
 fax: (509) 659-4109  
 email: [karenp@co.adams.wa.us](mailto:karenp@co.adams.wa.us)



website: <http://www.co.adams.wa.us/departments/>

TECHNICAL NOTES: TOTAL DEATHS DUE TO TOBACCO USE ARE ESTIMATED AS 19% OF ALL DEATHS (MCGINNIS AND FOEGE 1993); "DUMBELL" BARS AROUND DATA POINTS FOR SURVEY DATA REPRESENT 95% CONFIDENCE INTERVALS – THE TRUE VALUE MAY BE ANYWHERE WITHIN THE BARS, BUT MAY NOT BE EXACTLY AT THE DATA POINT.

ADDITIONAL DOCUMENTATION IN APPENDIX B: [http://www.doh.wa.gov/Tobacco/Data/County\\_Profiles/Profiles\\_2002/AppndxB.pdf](http://www.doh.wa.gov/Tobacco/Data/County_Profiles/Profiles_2002/AppndxB.pdf)

MAPS OF COUNTY RANKINGS: : [http://www.doh.wa.gov/Tobacco/Data/County\\_Profiles/Profiles\\_2002/AppndxC.pdf](http://www.doh.wa.gov/Tobacco/Data/County_Profiles/Profiles_2002/AppndxC.pdf)

REFERENCES: [http://www.doh.wa.gov/Tobacco/Data/County\\_Profiles/Profiles\\_2002/AppndxD.pdf](http://www.doh.wa.gov/Tobacco/Data/County_Profiles/Profiles_2002/AppndxD.pdf)